

Rennick-Egglestone, S. et al. (2019) Mental health recovery narratives and their impact on recipients: systematic review and narrative synthesis. *Canadian Journal of Psychiatry*, (doi:[10.1177/0706743719846108](https://doi.org/10.1177/0706743719846108)).

This is the author's final accepted version.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/183679/>

Deposited on: 09 April 2019

Mental health recovery narratives and their impact on recipients: systematic review and narrative synthesis

Stefan Rennick-Egglestone ^{a*}, Kate Morgan ^a, Joy Llewellyn-Beardsley ^a, Amy Ramsay ^b, Rose McGranahan ^c, Steve Gillard ^d, Ada Hui ^a, Fiona Ng ^a, Justine Schneider ^e, Susie Booth ^f, Vanessa Pinfold ^g, Larry Davidson ^h, Donna Franklin ^f, Simon Bradstreet ⁱ, Simone Arbour ⁱ, Mike Slade ^a

^a School of Health Sciences, Institute of Mental Health, University of Nottingham

^b Health Service and Population Research Department, Institute of Psychiatry, Psychology and Neuroscience, King's College London

^c Unit of Social and Community Psychiatry, Blizard Institute, Barts and the London School of Medicine and Dentistry, Queen Mary University of London

^d Population Health Research Institute, St. George's University of London

^e School of Sociology and Social Policy, University of Nottingham

^f NEON Lived Experience Advisory Panel

^g McPin Foundation

^h Yale School of Medicine, Yale University

ⁱ Institute of Health and Wellbeing, University of Glasgow

^j Ontario Shores Centre for Mental Health Science

* Rennick-Egglestone is the corresponding author.

Institute of Mental Health, Triumph Road, Nottingham, NG7 2TU, United Kingdom.

stefan.egglestone@nottingham.ac.uk +44 115 30926

Abstract

Objective. Mental health recovery narratives are often shared in peer support work and anti-stigma campaigns. Internet technology provides access to an almost unlimited number of narratives, and yet little is known about how they impact on recipients. The aim of this study was to develop a conceptual framework characterising the impact of recovery narratives on recipients.

Method. A systematic review of evidence about the impact of mental health recovery narratives was conducted. Searches used electronic databases (n=9), reference tracking, hand-searching of selected journals (n=2), grey literature searching and expert consultation (n=7). A conceptual framework was generated through a thematic analysis of included papers, augmented by consultation with a Lived Experience Advisory Panel.

Results. 8,137 articles were screened. Five papers were included. Forms of impact were: Connectedness; Understanding of recovery; Reduction in stigma; Validation of personal experience; Affective responses. Behavioural responses. Impact is moderated by characteristics of the recipient, context and narrative. Increases in eating disorder behaviours was identified as a harmful response specific to recipients with eating disorders.

Conclusions. Mental health recovery narratives can promote recovery. Recovery narratives might be useful for clients with limited access to peers, and in on-line interventions targeted at reducing social isolation in rural or remote locations, but support is needed for the processing of the strong emotions which can arise. Caution is needed for use with specific clinical populations.

Protocol registration: Prospero-CRD42018090923.

Keywords. Mental health; Recovery narratives; Recovery stories; Memoirs; Impact

Introduction

Mental health recovery narratives are first-person accounts of recovery from mental health problems which refer to events or actions over a period of time.¹ They are referred to as 'recovery narratives' throughout the remainder of this paper, whilst recognising that this term is used elsewhere in healthcare research and practice, e.g. in narratives of recovery post-stroke.²

Recovery narratives can be live (given in the context of a real-world or online relationship) or recorded (presented in invariant text, video or audio). Live recovery narratives feature in peer work, where professional training typically encourages the selective disclosure of personal experiences, as part of an ongoing relationship with a client. A USA national survey has identified helping others through the narrating of recovery narratives as a feature of the work of peer specialists,³ and Davidson, et al. ⁴ have argued that the disclosure by a peer worker of their own transition to a "hero of their own self-journey" (p.124) can instil hope in others.

Substantial numbers of recovery narratives are publicly available, distributed through books^{5,6}, on-line collections⁷ and digital media hosting services.⁸ Creating narratives can provide benefits for narrators,⁹ who might be motivated by sending messages of "hope, courage and survival" (p. 68),¹⁰ a form of indirect emotional support provision.¹¹ Campaigns which aim to reduce stigma^{12,13}, such as Bell Let's Talk¹⁴, have used recorded recovery narratives¹⁵ as a mechanism for creating social contact^{16,17} between people with experience of mental health problems and others. Narratives used for anti-stigma purposes can also have a beneficial impact on help-seeking behaviour,¹⁸ and organisations such as Here to Help¹⁹ and the Scottish Recovery Network²⁰ have created on-line collections with the explicit intent of supporting recovery in recipients. These might be seen as a specific initiatives within a larger effort to incorporate information and communication technologies into mental health practice, motivated by challenges

such as limited access to in-person mental health treatment in rural^{21,22} and First Nations²³ communities, and for people with social anxiety disorders.²⁴

Despite widespread interest, a knowledge gap exists about how live and recorded recovery narratives make an impact. No systematic review has been conducted on this topic. A realist review considered the broader question of how sharing experiences of health conditions on the Internet might affect the health of those sharing and receiving material; it identified mechanisms such as (1) the recipient finding useful information in the experiences of others with similar health difficulties, or (2) feeling supported through an understanding that others have encountered and overcome related challenges.²⁵ This review drew on a range of empirical and non-empirical sources. Its authors note a need to strengthen the scientific evidence base, given an “*explosion of web resources that feature experiences posted by patients themselves*” (p. 219). The authors note that receiving experiences had the potential for positive and negative impact.

Other studies have identified specific harms that can be associated with receiving on-line health material, especially in the case of conditions that are isolating or stigmatised.²⁶ Examples include “pro-ana” websites, which enable on-line inter-person interaction that sustains anorexic behaviours and beliefs.²⁶ An experimental study of 235 female undergraduates showed that viewing a pro-anorexia website increased negative affect and lowered social self-esteem relative to a comparison website.²⁷ A qualitative study demonstrated that pro-anorexia sites can sustain anorexic behaviour by enabling the sharing of hints and tips.²⁸ Neither study was specific to narratives, but included websites utilising narrative material.

Mental health recovery narratives are becoming more widely available, so an evidence-based understanding of how they make an impact on recipients is needed, especially given known harms associated with on-line health material. The aim of this review was to develop a conceptual framework characterising the impact of live or recorded recovery narratives on recipients.

Method

A systematic review of documents providing empirical evidence for the impact of recovery narratives on recipients was conducted. The systematic review protocol was registered:

http://www.crd.york.ac.uk/PROSPERO/display_record.php?ID=CRD42018090923.

Recovery narratives were defined as first-person non-fiction accounts of recovery from mental health problems, including (1) elements of adversity/struggle and (2) self-defined strengths/successes/survival, and which (3) refer to events or actions over a period of time.¹

“Receiving a narrative” was defined as viewing, reading or listening to someone else’s recovery narrative. “Impact” was defined as the processes by which receiving recovery narratives causes benefits or harms to the recipient and the outcomes generated by these.

Inclusion criteria: (1) presents empirical research, with at least five participants as recipients of recovery narratives; (2) presents evidence relating to the impact of recovery narratives on participants, including evidence of no impact; (3) draws on non-fictional narratives with a clear mental health and recovery component; (4) the full text of the publication was available in English.

Exclusion criteria: (1) narratives were told second-hand, e.g. by a healthcare professional or carer (2) narrative were a component of a complex intervention and it was not clear whether the impact was due to the narrative or to other components of the intervention. The latter was included due to scoping searches locating papers in which recovery narratives were just one part of a multi-component intervention evaluated as a whole, and where the impact of individual components could not be disambiguated. An example was a two-component anti-stigma campaign²⁹ which utilised recovery narratives and educational workshops.

Search strategy

Electronic databases

Research into recovery narratives is multidisciplinary, and hence searches were conducted across a broad range of publication databases. Selection was informed by scoping searches. Selected databases were: AMED, EMBASE, MEDLINE, PSYCINFO (all via OVID). ASSIA, MLA, PILOTS, CINAHL, Web of Science. To avoid a risk of omitting relevant material, scoping searches identified a range of terms that might be loosely synonymous for “recovery narrative”, including “memoirs” and “autobiographies”.

Database search terms were piloted and then defined against AMED (via OVID), to generate a reference search strategy presented in the digital supplement. This was specialised to each database. Publications were included from inception until 31 August 2018.

Publications were filtered by title and abstract by KM, with concordance checking by SRE on a sample of 10% of titles and 20% of abstracts. Concordance was 96% for title filtering, and 95% for abstract filtering. Remaining papers were retrieved. Inclusion was decided by KM and SRE, with disagreements resolved by discussion with other authors. If a candidate publication did not use the term “recovery narrative”, it was included if definitive evidence could be assembled from the abstract or full text that the narratives it considered were recovery narratives. This involved inspecting specific narratives listed in publications for mental health and recovery content.

Hand searching of journals

The two journals containing papers included through database searches (*Advances in Eating Disorders: Theory, Research and Practice* and *Journal of Nervous and Mental Disease*) were hand-searched from inception to 31 August 2018.

Grey literature searching

Google and Google Scholar were searched using the terms “Impact of mental health recovery narratives” and “Impact of mental health recovery stories”. The first 100 hits were hand-searched.

Expert consultation

Seven researchers with expertise in mental health recovery were sent the list of included papers and asked to identify additional candidates for inclusion.

Citation tracking

References lists of all included papers and of a relevant review¹³, were hand-searched. Forward citation tracking of included papers and this review was conducted using Google Scholar.

Data abstraction

A data abstraction table (DAT) was designed and piloted, and is presented in the digital supplement. The DAT presents details of each included study: country of lead author, design of study, characteristics of narratives, characteristics of participants, time of impact assessment, social interaction around the narrative. It also presents knowledge about the impact of recovery narratives generated by each study, grouped by categories refined through the piloting process: form of impact, consequences of impact, facilitators to impact and barriers to impact. Included qualitative papers presented rich narrative descriptions of impact. This meant that the work of data abstraction involved producing short textual summaries of narrative descriptions. Summaries were agreed by the two lead authors, and lead by the language of the source publication.

Quality assessment

Included studies were critically appraised. Results are included in the Data Abstraction Table. The Critical Appraisal Skills Programme Qualitative Checklist³⁰ was used for qualitative studies, using established thresholds to indicate high, moderate or low rated quality³¹. Quantitative studies were assessed using the 15-element McMaster Critical Review Form for Quantitative Studies³², with bespoke labels: high (13-15), moderate (11-12), low (0-10).

Narrative synthesis

A narrative synthesis was conducted following an established methodology³³. This consisted of (1) a preliminary synthesis of abstracted material on forms, consequences, facilitators and barriers to impact (2) refinement of the synthesis through discussion in a multi-disciplinary research team (3) refinement of the synthesis through consultation with a Lived Experience Advisory Panel and (4) sub-group analysis. The goal of the synthesis process was to generate a framework describing broad forms of impact, but which retained essential details of how impact occurred as presented in included papers.

The preliminary synthesis began by classifying knowledge about impact presented in the DAT as (1) transdiagnostic or (2) diagnostically specific. Example: “Feeling connected to the narrator” was classified as transdiagnostic, whilst “greater effort put into avoidance of ED behaviours” was classified as specific to an eating disorder diagnosis. Classifications were by SRE, and were reviewed by clinically-qualified authors.

A preliminary synthesis was generated by SRE, who conducted an inductive thematic analysis on items classified as transdiagnostic. Items were grouped into named themes, which were selected (1) to describe a meaningful form of impact (2) for orthogonality. Similar items grouped within a

theme were combined. To raise quality, the preliminary synthesis was discussed in a team comprising analysts from multi-professional backgrounds (mental health nursing, social psychiatry, clinical psychology, community development, design research) and including qualitative expertise. Closely related items were further combined, and names of themes and items were refined for clarity. 'Moderators' was introduced to encompass facilitators and barriers.

The updated synthesis was presented to a Lived Experience Advisory Panel (LEAP), comprising ten members with prior experience of mental health problems and of recovery. Three had published their own recovery narrative. A final version was produced based on their input. The principle change was identification of items to add existing themes to add depth. These items have been labelled as LEAP contributions to distinguish them from items sourced from included publications. During this consultation, LEAP members raised questions relevant to future research, and these are considered in the Discussion.

Items labelled as diagnostically specific to a clinical population were examined. To document forms of impact specific to that population, processes, outcomes and moderators were tabulated.

Results

The PROSPERO flow diagram for this review is shown in Figure 1.

Insert Figure 1 here

Five papers were included (#1,#2,#3,#4,#5). Featured narratives were characterised as “recovery stories” (#1,#5), “memoirs” (#2, #3) and “lived experience videos” (#4). All narratives present in #2, #3 and #4 were inspected, and found to meet the study definition of a recovery narrative.

Key characteristics of included studies are in Table 1. Further detail is provided in the DAT.

Insert Table 1 here

All included papers were studies of the impact of recorded recovery narratives, and were of moderate or high quality. The single included quantitative study, #2, failed to find evidence for impact of an eating disorder memoir on participants who were female undergraduates, and not selected for membership of a clinical population. As this paper provided no evidence for impact, facilitators or barriers, then it has not informed the narrative synthesis.

The four included qualitative studies (#1, #3, #4, #5) were conducted recently. Collectively, these papers included 93 participants, 70 female and 23 male, and hence there is a clear gender bias across included qualitative research. Papers #1, #3 and #4 were conducted by research teams based in Australia, but with no overlap in authorship. #1, #3 and #4 note a lack of existing empirical evidence about the impact of recovery narratives on recipients.

Specific items were identified that provided insights into the impact of mental health recovery narratives on individuals experiencing eating disorders, and a subgroup analysis was conducted on these. Despite the inclusion of a paper whose participants had been diagnosed with a psychotic condition, no knowledge about impact specific to psychosis was identified.

Framework for transdiagnostic impact

Table 2 presents a framework describing transdiagnostic impact, synthesised from forms and consequences identified in the DAT. This table includes the six themes identified through thematic analysis, which are our proposed forms of impact, and the items grouped within each theme, which provide more specific detail about how they occur.

Insert Table 2 here

Impact described in this table has not been categorised as helpful or harmful, due to the possible influence of contextual factors, e.g. “initiating personal disclosure to a researcher” might be

helpful if the researcher has the required competence to engage, but harmful if not.

Connectedness, Understanding, Reduction in Stigma and Validation might be thought of as more cognitive forms of impact, alongside the broad range of more affective responses listed in table 2, and a limited range of behavioural responses.

Table 3 presents moderators identified through the synthesis, identified from facilitators and barriers listed in the DAT.

Insert Table 3 here

Items included in the categories of recipient and narrator/narrative characteristics appeared to facilitate or block a recipient in forming a connection with the narrator or narrative, suggesting an important role for connection forming in enabling impact to occur.

Forms of impact specific to eating disorders

Table 4 summarises processes and impact identified as specific to eating disorders.

Insert Table 4 here

All identified processes can generate harmful behavioural responses, though for the first process this only occurred if the recipient did not identify as being “in recovery” at the time of receiving the narrative; if the participant did identify as being “in recovery” or “recovered” a helpful behavioural outcome was generated. These terms were participant-defined in the source paper, e.g. there was no usage of a standardised measure to assess recovery stage.

Paper 3 identifies a *narrator/narrative characteristic* that facilitates harmful behavioural outcomes - the inclusion of particular details of eating disorder behaviours engaged in by the narrator, such as dangerously low daily calorie counts or specific patterns of disordered eating.

Discussion

This review has presented a conceptual framework consisting of nine superordinate themes, comprising six types of impact and three moderators of impact. A sub-group analysis identified types of impact specific to eating disorders. The inclusion of only five papers, despite widespread interest in the use of recovery narratives, suggests a knowledge gap which might be filled through further empirical work, and identification of this knowledge gap is a key contribution of this paper. These findings will inform a future trial (ISRCTN11152837).

Relationship to existing frameworks

A prior review²⁵ has considered the impacts of sharing and receiving health material online, and identified four domains in which positive and negative impact can occur for recipients: (1) finding information (2) feeling supported (3) experiencing health services (4) affecting behaviour. *Feeling supported* has similarities with *Validation of personal experiences*, whilst *finding information* has similarities with *Understanding of mental illness and how to recover* (with the latter being more specific to recovery narratives). Impact on behaviour occurs in both.

Our review has a stronger emphasis on factors that moderate impact, only briefly mentioned in the previous review. It also emphasises forms of impact distinctive to recovery narratives, such as connection with and empathy for the narrator. Our review does not consider issues around unbalanced or misleading material, considered extensively in the previous review. Learning that an influential narrative might not be authentic can lead to anger in recipients³⁴, and LEAP members wondered whether recipients could be harmed by deliberately misleading narratives.

Some on-line collections of recovery narratives are predicated on a belief that receiving narratives can encourage personal recovery. A systematic review has identified Connectedness, Hope;

Identity; Meaning in life; and Empowerment (CHIME) as processes that underpin personal recovery³⁵. Elements of all five CHIME process are present in our conceptual framework. Connectedness appears in both frameworks. The CHIME notion of Hope appears under Affective responses, whilst the CHIME notion of Identity incorporates “Overcoming stigma”, present as an impact in our conceptual framework. This provides some validation that receiving recovery narratives can promote recovery. The inclusion of “Reduction in stigma” as an outcome provides validation for the use of recovery narratives in anti-stigma campaigns.

Strengths and limitations

A strength of the review is the identification, through scoping searches, of a broad range of terms that might be synonymous of “recovery narrative”, important as no included paper used this term directly. Other strengths include the use of a wide range of publication databases and grey literature sources, double rating at each stage of sifting to maximise reliability, and the use of an established methodology for narrative synthesis. Limitations include a narrow focus on mental health recovery narratives, given that the impact of narratives produced by other clinical populations may also be relevant in mental health practice, the non-use of some potential sources such as conference proceedings, and the restriction of the search strategy to English, meaning that some French-language publications could have been excluded.

Implications for clinical practice and mental health interventions

Should recovery narratives be used as a resource in clinical practice and mental health interventions? The general clinical evidence is supportive for the former. A survey of 138 psychologists in the USA found a belief that mental health biographies can be helpful in therapy sessions³⁶, and a review of case reports has demonstrated mechanisms for integrating

autobiographies into psychotherapy sessions³⁷. A range of publications have shown acceptability of bibliotherapy and self-help texts as a form of treatment across a range of conditions^{38,39}. Biographical material used in these studies was typically in the form of recovery narratives.

The ready availability of mental health recovery narratives on-line may support usage with younger people, who are known to be heavy users of on-line health information.⁴⁰ This is important given recent arguments that youth mental health should be a priority area in Canada,⁴¹ and known challenges around extremely high suicide rates in Canadian Aboriginal youth.⁴² Best practice guidelines on Canadian youth suicide reduction⁴³ include ensuring accessibility of services in rural and remote locations and enhancing engagement with marginalised individuals. Interventions drawing on on-line material are known to have particular relevance to areas with low population densities²², where access to both mental health care and to peers with experience of similar mental health problems may be difficult otherwise⁴⁴. Although limited, internet access in rural and remote areas of Canada is improving,⁴⁵ including through enhancements to cellular networks,⁴⁶ and initiatives such as the Indigenous Connectivity Summit.⁴⁷

Our review highlights potential mental health uses and considerations that are relatively specific to recovery narratives. People experiencing high anticipated discrimination may benefit from stigma reduction by encountering role models through narratives, who may not be available in everyday life. People experiencing high levels of social isolation, a known risk factor for youth suicide⁴⁸, may experience enhanced connectedness and more validation of personal experiences. People who are early in their recovery journey may encounter relevant knowledge about how to recover. Knowledge presented in recovery narratives might enable people to cope more readily with the challenges of recovery, with coping being one source of resilience⁴⁹.

The review found that recovery narratives can engender strong affective responses, sometimes through empathy to the narrator, including emotions perceived as positive (hopefulness, gratitude

for life) and others that are more distressing. LEAP members queried whether secondary traumatisation might be one outcome of excessive exposure to distressing elements of recovery narratives, potentially through mechanisms similar to those known in relation to mental healthcare providers⁵⁰. When available, clinicians should monitor and support processing of emotional responses by clients, given that empathy is known to be significant risk factor for secondary traumatisation⁵¹. Alternative approaches are needed when recovery narratives are used in on-line mental health interventions. These might include moderated discussion forums or on-line support groups, which can offer emotional and instrumental¹¹ support to members.⁵²

Our review found no material specific to psychosis, despite inclusion of a paper with participants diagnosed with a psychotic condition, and hence does not provide any argument against the use of recovery narratives in the psychosis treatment– in the context of a debate around the appropriateness of using on-line material in the treatment of psychotic disorders⁵³.

A lack of connectedness to the narrator of recovery narrative is identified as a significant barrier to impact in Table 3, and is also indicated as a potential cause of distress. This suggests a need for further research to identify factors that might disrupt connection, of broader importance given the perceived importance of personal connection in peer support work. It also highlights the need for clinicians to attend to a client's capacity to connect as part of the decision-making process around whether to use introduce recovery narratives, and when making a selection.

In keeping with prior research into on-line material and eating disorders, this review supports an understanding of harms that can be caused by engaging with recovery narratives, through processes linked to the elements of recovery narrative presenting adversity and struggle. LEAP members queried whether similar mechanisms might be seen in consumers with prior experience of self-harm, consistent with prior research which suggests that on-line material can in fact trigger instances of self-harm⁵⁴. In keeping with Whitlock, et al. ⁵⁵, we would argue that regular

assessment of narrative usage might enable the amelioration of harms. Alternative mechanisms are needed in the case of on-line interventions.

The findings of our review accord with prior work which suggests that the dissemination of specific details of harmful behaviours in on-line material is one mechanism of harm⁵⁶. This might impact on processes for selecting recovery narratives for use in mental health practice, through tactics such as avoiding recovery narratives that include specific details of harmful behaviours.

Conclusions

Mental health recovery narratives can impact on cognition, affect and behaviour, and this impact is moderated by a range of factors. Mental health recovery narratives can promote recovery in recipients, but care might be needed for use with specific clinical populations.

Data Access

The complete Data Abstraction Table for this work is included as an on-line supplement.

Acknowledgements

There are no acknowledgements in addition to the funding acknowledgements below.

Conflicts of Interest Disclosure

The Authors declares that there is no conflict of interest.

Funding

This article is independent research funded by the NIHR under its Programme Grants for Applied Research Programme (Programme Grants for Applied Research, Personal experience as a recovery resource in psychosis: Narrative Experiences ONline (NEON) Programme, RP-PG-

0615-20016). Mike Slade acknowledges the support of Center for Mental Health and Substance Abuse, University of South-Eastern Norway and the NIHR Nottingham Biomedical Research Centre. The views expressed are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care.

References

1. Llewellyn-Beardsley J, Rennick-Egglestone S, Callard F, Crawford P, Farkas M, Hui A, Manley D, McGranahan R, Pollock K, Ramsay A and others. Characteristics of mental health recovery narratives: systematic review and narrative synthesis (in press). *PLOS One* 2019.
2. Faircloth CA, Boylstein C, Rittman M, Young ME, Gubrium J. Sudden illness and biographical flow in narratives of stroke recovery. *Sociology of Health & Illness* 2004;26(2):242-261.
3. Cronise R, Teixeira C, Rogers ES, Harrington S. The peer support workforce: Results of a national survey. *Psychiatric Rehabilitation Journal* 2016;39(3):211.
4. Davidson L, Bellamy C, Guy K, Miller R. Peer support among persons with severe mental illnesses: a review of evidence and experience. *World Psychiatry* 2012;11(2):123-128.
5. Cordle H, Carsen J, Richards P, Fradgley J. *Psychosis: Stories of Recovery and Hope*. Quay Books; 2010.
6. Chandler R, Hayward M. *Voicing psychotic experiences: A reconsideration of recovery and diversity*. OLM-Pavilion; 2009.
7. McDonaugh T, Crane L. The Schizophrenia Oral History Project. <<https://schizophreniaoralhistories.com/>>. Archived at: <http://www.webcitation.org/76NW2gcAC>>. Accessed 26th February 2019.
8. Poolay A. 2015. The human element of recovery from mental illness and addiction. In TedX talk at Michigan State University. <https://www.youtube.com/watch?v=-m_A8ST_18s>.
9. Pennebaker JW, Seagal JD. Forming a story: The health benefits of narrative. *Journal of Clinical Psychology* 1999;55(10):1243-1254.
10. Kirkpatrick H, Byrne C. A narrative inquiry: Moving on from homelessness for individuals with a major mental illness. *Journal of Psychiatric and Mental Health Nursing* 2009;16(1):68-75.
11. Morelli SA, Lee IA, Arnn ME, Zaki J. Emotional and instrumental support provision interact to predict well-being. *Emotion* 2015;15(4):484.
12. Thornicroft G, Mehta N, Clement S, Evans-Lacko S, Doherty M, Rose D, Koschorke M, Shidhaye R, O'Reilly C, Henderson C. Evidence for effective interventions to reduce mental-health-related stigma and discrimination. *The Lancet* 2016;387(10023):1123-1132.
13. Evans-Lacko S, London J, Japhet S, Rüsch N, Flach C, Corker E, Henderson C, Thornicroft G. Mass social contact interventions and their effect on mental health related stigma and intended discrimination. *BMC Public Health* 2012;12(1):489.
14. Bell Let's Talk. Campaign website. <<https://letstalk.bell.ca/>>. Accessed 16th March 2019.

15. Lemoine P. Recovery narrative. <<https://letstalk.bell.ca/en/our-initiatives/team/patricia-lemoine>>. Accessed 16th March 2019.
16. Allport GW, Clark K, Pettigrew T. The nature of prejudice. Addison-Wesley; 1979.
17. Corrigan PW, River LP, Lundin RK, Penn DL, Uphoff-Wasowski K, Campion J, Mathisen J, Gagnon C, Bergman M, Goldstein H. Three strategies for changing attributions about severe mental illness. *Schizophrenia Bulletin* 2001;27(2):187-195.
18. Henderson C, Robinson E, Evans-Lacko S, Thornicroft G. Relationships between anti-stigma programme awareness, disclosure comfort and intended help-seeking regarding a mental health problem. *The British Journal of Psychiatry* 2017;211(5):316-322.
19. Here to Help. Personal stories. <<http://www.heretohelp.bc.ca/personal-stories>>. Archived at: <http://www.webcitation.org/76NWHi7mz>>. Accessed 26th February 2019.
20. Bradstreet S. 2009. Scottish Recovery Network Story Sharing Guidelines. <<https://www.scottishrecoveryconsortium.org/assets/files/SRN%20Story%20sharing%20guide%202009%20-%20new%20logo.pdf>>. Archived at: <http://www.webcitation.org/76Ndhnk72>>. Accessed 26th February 2019.
21. Cassidy L. Online communities of practice to support collaborative mental health practice in rural areas. *Issues in Mental Health Nursing* 2011;32(2):98-107.
22. Griffiths KM, Christensen H. Internet based mental health programs: A powerful tool in the rural medical kit. *Australian Journal of Rural Health* 2007;15(2):81-87.
23. Gibson K, Simms D, O'Donnell S, Molyneaux H. Clinicians' attitudes toward the use of information and communication technologies for mental health services in remote and rural areas. *Proceedings of Canadian Society of Telehealth Conference*; 2009; Vancouver, British Columbia.
24. Andersson G, Titov N. Advantages and limitations of Internet-based interventions for common mental disorders. *World Psychiatry* 2014;13(1):4-11.
25. Ziebland S, Wyke S. Health and illness in a connected world: how might sharing experiences on the internet affect people's health? *The Milbank Quarterly* 2012;90(2):219-249.
26. Norris ML, Boydell KM, Pinhas L, Katzman DK. Ana and the Internet: A review of pro-anorexia websites. *International Journal of Eating Disorders* 2006;39(6):443-447.
27. Bardone-Cone AM, Cass KM. What does viewing a pro-anorexia website do? An experimental examination of website exposure and moderating effects. *International Journal of Eating Disorders* 2007;40(6):537-548.
28. Mulveen R, Hepworth J. An interpretative phenomenological analysis of participation in a pro-anorexia internet site and its relationship with disordered eating. *Journal of Health Psychology* 2006;11(2):283-296.
29. Pinfold V, Thornicroft G, Huxley P, Farmer P. Active ingredients in anti-stigma programmes in mental health. *International Review of Psychiatry* 2005;17(2):123-131.
30. Singh J. Critical appraisal skills programme. *Journal of Pharmacology and Pharmacotherapeutics* 2013;4(1):76.
31. Butler A, Hall H, Copnell B. A guide to writing a qualitative systematic review protocol to enhance evidence-based practice in nursing and health care. *Worldviews on Evidence-Based Nursing* 2016;13(3):241-249.
32. Law M, Stewart, D., Pollock, N., Letts, L. Bosch, J., & Westmorland, M. 2018. Critical Review Form – Quantitative Studies. <https://www.unisa.edu.au/Global/Health/Sansom/Documents/iCAHE/CATs/McMasters_Quantitative%20review.pdf>. Accessed 2nd November 2018.
33. Popay J, Roberts H, Sowden A, Petticrew M, Arai L, Rodgers M, Britten N, Roen K, Duffy S. Guidance on the conduct of narrative synthesis in systematic reviews. A product from the ESRC methods programme Version 2006;1:b92.

34. Rennick-Egglestone S, Knowles S, Toms G, Bee P, Lovell K, Bower P. Health Technologies' In the Wild': Experiences of Engagement with Computerised CBT. Proceedings of 2016 CHI Conference on Human Factors in Computing Systems; 2016. ACM. p 2124-2135.
35. Leamy M, Bird V, Le Boutillier C, Williams J, Slade M. Conceptual framework for personal recovery in mental health: systematic review and narrative synthesis. *The British Journal of Psychiatry* 2011;199(6):445-452.
36. Clifford JS, Norcross JC, Sommer R. Autobiographies of mental health clients: Psychologists' uses and recommendations. *Professional Psychology: Research and Practice* 1999;30(1):56.
37. Sommer R. The use of autobiography in psychotherapy. *Journal of Clinical Psychology* 2003;59(2):197-205.
38. Kilfedder C, Power K, Karatzias T, McCafferty A, Niven K, Chouliara Z, Galloway L, Sharp S. A randomized trial of face-to-face counselling versus telephone counselling versus bibliotherapy for occupational stress. *Psychology and Psychotherapy: Theory, Research and Practice* 2010;83(3):223-242.
39. Brenes GA, McCall WV, Williamson JD, Stanley MA. Feasibility and acceptability of bibliotherapy and telephone sessions for the treatment of late-life anxiety disorders. *Clinical Gerontologist* 2009;33(1):62-68.
40. Burns JM, Davenport TA, Durkin LA, Luscombe GM, Hickie IB. The internet as a setting for mental health service utilisation by young people. *Medical Journal of Australia* 2010;192(11):S22.
41. Malla A, Shah J, Iyer S, Boksa P, Joober R, Andersson N, Lal S, Fuhrer R. Youth Mental Health Should Be a Top Priority for Health Care in Canada. *The Canadian Journal of Psychiatry* 2018;63(4):216-222.
42. Health Canada. National Aboriginal Youth Suicide Prevention Strategy (NAYSPS): Program Framework. 2013.
43. Kirmayer LJ, Fraser S-L, Fauras V, Whitley R. Current approaches to aboriginal youth suicide prevention. Cultural and Mental Health Research Unit working paper 2009;14.
44. Serhal E, Crawford A, Cheng J, Kurdyak P. Implementation and Utilisation of Telepsychiatry in Ontario: A Population-Based Study. *The Canadian Journal of Psychiatry* 2017;62(10):716-725.
45. Ruimy D. Broadband connectivity in rural Canada: overcoming the digital divide. Standing Committee on Industry, Science and Technology;; 2018.
46. Coop A. 2018. Deloitte: 25 per cent of Canadians will get internet from cellular mobile networks this year. <<https://www.itbusiness.ca/news/deloitte-25-per-cent-of-canadians-will-get-internet-from-cellular-mobile-networks-this-year/98187>>.
47. Indigenous Connectivity Summit. 2017. Summit website. <<https://www.internetsociety.org/events/indigenous-connectivity-summit/2017/>>.
48. King CA, Merchant CR. Social and interpersonal factors relating to adolescent suicidality: a review of the literature. *Archives of Suicide Research* 2008;12(3):181-196.
49. Pai N, Vella S-L. Can one spring back from psychosis? The role of resilience in serious mental illness. *The Australian and New Zealand Journal of Psychiatry* 2018;4867418802900.
50. Zimering R, Munroe J, Gulliver S. Secondary traumatization in mental health care providers. *Psychiatric Times* 2003;20(4).
51. Crumpei I, Dafinoiu I. The relation of clinical empathy to secondary traumatic stress. *Procedia-Social and Behavioral Sciences* 2012;33:438-442.
52. Wang Y-C, Kraut R, Levine JM. To stay or leave?: the relationship of emotional and informational support to commitment in online health support groups. Proceedings of

- ACM 2012 conference on Computer Supported Cooperative Work; 2012. ACM. p 833-842.
53. Thompson A, Gleeson J, Alvarez-Jimenez M. Should we be using digital technologies in the treatment of psychotic disorders? *Australian & New Zealand Journal of Psychiatry* 2018;52(3):225-226.
 54. Jacob N, Evans R, Scourfield J. The influence of online images on self-harm: a qualitative study of young people aged 16–24. *Journal of Adolescence* 2017;60:140-147.
 55. Whitlock J, Lader W, Conterio K. The internet and self-injury: what psychotherapists should know. *Journal of Clinical Psychology* 2007;63(11):1135-1143.
 56. Bell J. Harmful or helpful? The role of the internet in self-harming and suicidal behaviour in young people. *Mental Health Review Journal* 2014;19(1):61-71.
 57. Sheens E, Rhodes P, Dawson L. Encountering anorexia: challenging stigma with recovery stories. *Advances in Eating Disorders* 2016;4(3):315-322.
 58. Thomas JJ, Judge AM, Brownell KD, Vartanian LR. Evaluating the effects of eating disorder memoirs on readers' eating attitudes and behaviors. *International Journal of Eating Disorders* 2006;39(5):418-425.
 59. Shaw L-K, Homewood J. The effect of eating disorder memoirs in individuals with self-identified eating pathologies. *The Journal of Nervous and Mental Disease* 2015;203(8):591-595.
 60. Williams A, Fossey E, Farhall J, Foley F, Thomas N. Recovery After Psychosis: Qualitative Study of Service User Experiences of Lived Experience Videos on a Recovery-Oriented Website. *JMIR Mental Health* 2018;5(2).
 61. Nurser KP, Rushworth I, Shakespeare T, Williams D. Personal storytelling in mental health recovery. *Mental Health Review Journal* 2018;23(1):25-36.

Figures and tables

Figure 1: Flow diagram




Table 1 Summary of included publications (n=5)

#	Year	Design	Ref	Description	Time of impact assessment	Quality rating
1	2016	Qual (n=25)	⁵⁷	Thematic analysis of semi-structured interviews with first year psychology students who listened to an Anorexia Nervosa recovery narrative.	Immediately after hearing the story	Moderate
2	2006	Quant (n=54)	⁵⁸	Study of the impact of a written eating disorder recovery narrative on female undergraduate Psychology students. Impact assessed using the Eating Attitudes Test (EAT-26), the Eating Disorders Inventory (EDI) Drive for Thinness subscale, a measure of perceived ED symptom prevalence, and an Implicit Association Test (IAT) measuring associations between anorexia and glamour/danger	Within one week of reading the story	Moderate
3	2015	Qual (n=24)	⁵⁹	Qualitative analysis of questionnaires completed by participants who consumed eating disorder recovery narratives as part of their own recovery	Undefined	Moderate

4	2018	Qual (n=36)	⁶⁰	Interviews with participants who have been diagnosed with a psychotic disorder, and who have engaged with used video-based recovery narratives presented on a website as part of their recovery.	At least three months after initial contact with website	High
5	2018	Qual (n=8)	⁶¹	Interpretive Phenomenological Analysis of interviews with attendees of a Telling My Story course.	Within one year of taking the course	High

Table 2: Impact and associated processes of recovery narratives on recipients

Impact	Associated processes	Source
Connectedness to the narrator or to others	The recipient feels connected to the narrator of a recovery narrative	1,4
	The narrative provides companionship at times of social isolation	3
	The narrative helps the recipient to feel that they belong	4
Understanding of mental illness and how to recover	The recipient's understanding of the severity, intensity and complexity of the mental health condition experienced by the narrator is deepened	1
	The recipient's understanding of how personal recovery might happen is deepened	1,4
	The recipient identifies personal behaviours that they wish to reconsider	4
	The recipient learns from a narrator who has experienced more profound difficulties than them	LEAP
Reduction in stigma (including self-stigma)	The recipient experiences a reduction in their stigmatic views about the mental health condition described by the narrator	1
	The recipient feels less ashamed about having a mental illness	4
	The recipient feels reassured that others have experienced similar distress	1

Validation of self and of personal experiences	The recipient hears hidden views about illness, and recognises themselves in these	4
	The recipient recognises some of their deepest fears in the narrative	LEAP
	The recipient compares themselves to the narrator, and finds that they are doing well	5
Affective responses	The recipient feels empathy for the narrator, and this initiates strongly-felt emotional responses (feeling moved, sad, distressed, heartbroken, inspired, admiring)	1,4 LEAP
	The recipient feels an increased gratitude for positive aspects of their life	1
	The recipient feels uncomfortable due to recall of difficult memories	LEAP
	The recipient feels increased hope that recovery is possible (and this hope provides comfort at times of distress)	3,4
	The recipient feels angry or out of place, as they have experienced significant distress, and the problems of the narrator feel insignificant	LEAP, 5
	The recipient feels pressured, as the narrative is of recovery achievements that the recipient cannot match	LEAP

	The recipient feels compelled to make personal changes after hearing a recovery narrative from someone they could relate to	5
Behavioural responses	The recipient initiates personal disclosure to the researcher	1
	The recipient initiates deeper discussion with a support worker	4
	The recipient becomes more forceful when talking about medication with medical practitioners	4

Table 3: Moderators of impact

Moderator	Description	Facilitator or Barrier	Source
Recipient characteristics	Life circumstances and events in the narrative matched those of the recipient	Facilitator	3,4,5
	The recipient reports a short-term or long-term inability to connect with others	Barrier	1
	The recipient is not able to relate to the narrator's journey of recovery, leading to feelings of alienation and despair		3,4
	The recipient has strongly held personal beliefs that contradict with those of the narrator		1
	The recipient avoids some recovery narratives due to the content being excessively challenging		4
	The recipient has an empathic response to video-based recovery stories, making consumption tiring		4
	The recipient is not at the right stage of recovery for a particular narrative to be beneficial		LEAP
	The recipient feels significantly worse off than the narrators, and holds back from connection		5

Contextual characteristics	The recipient had a support worker who helps with use of the technology	Facilitator	4
	The recipient had a choice of physical location to consume narratives		4
	Lack of access to internet technology at home	Barrier	4
Narrator and narrative characteristics	The narrator presents specific details of their own behaviours or life events	Either	3
	The narrator appears to find it hard to express emotions or be authentic	Barrier	LEAP

Table 4: Processes and impact specific to eating disorders narratives

Process	Impact	Source
Receiving a narrative enhances the recipient's understanding of the possible dangers of eating-disorder behaviours	[Recipient self-identifies as “in recovery” or “recovered”] Greater effort put into avoidance of eating disorder behaviours [Recipient self-identifies as not in recovery] Increases in belief that others are worse off, and hence change not necessary	3
Receiving a narrative encourages emulation of eating disorder behaviours enacted by the narrator	Increase in eating disorder behaviours	3
Receiving a narrative triggers prior eating disorder behaviours matching those of the narrator	Increase in eating disorder behaviours	3
Receiving a narrative leads to unhelpful social comparison with narrator	Increase in eating disorder behaviours	3